

FOR CLL/SLL


CALQUENCE[®]
acalabrutinib 100 mg tablets

START CALQUENCE

The first and only BTKi to offer two all-oral, chemo-free options for previously untreated CLL/SLL patients: fixed duration with **CALQUENCE + venetoclax** (completed in just over a year*) and continuous with **CALQUENCE** alone

What you need to know about your treatment

*Designed to be completed as prescribed in 12.9 months (fourteen 28-day cycles) unless your doctor tells you to stop. Your doctor will monitor you while on treatment, and will stop treatment if your disease progresses or if you develop certain side effects.

BTKi=Bruton tyrosine kinase inhibitor; CLL=chronic lymphocytic leukemia; SLL=small lymphocytic lymphoma.



Not an actual
patient.

Important Safety Information About CALQUENCE[®] (acalabrutinib)

Before taking CALQUENCE, tell your healthcare provider about all of your medical conditions, including if you:

- have had recent surgery or plan to have surgery. Your healthcare provider may stop CALQUENCE for any planned medical, surgical, or dental procedure.
- have bleeding problems.
- have or had heart rhythm problems.
- have an infection.
- have or had liver problems, including hepatitis B virus (HBV) infection.
- are pregnant or plan to become pregnant. CALQUENCE may harm your unborn baby and cause problems during childbirth (dystocia).

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with CALQUENCE
- Use effective birth control (contraception) during treatment with CALQUENCE and for 1 week after the last dose of CALQUENCE
- are breastfeeding or plan to breastfeed. It is not known if CALQUENCE passes into your breast milk. Do not breastfeed during treatment with CALQUENCE and for 2 weeks after your last dose of CALQUENCE.

Please see additional Important Safety Information throughout and click here for full Prescribing Information, including Patient Information.

TWO ORAL-ONLY, CHEMO-FREE TREATMENT OPTIONS

Learn more about the treatment options available with CALQUENCE



Fixed-duration treatment with CALQUENCE + venetoclax.

CALQUENCE in combination with venetoclax **can be completed in just over a year*** for adults with previously untreated CLL/SLL.



Continuous treatment with CALQUENCE.

CALQUENCE alone is a continuous treatment for adults with previously untreated and previously treated CLL/SLL.

Follow your treatment plan as prescribed by your doctor.
See pages 3-5 for dosage and administration instructions.



SEE CLINICAL STUDY RESULTS

*Designed to be completed as prescribed in 12.9 months (fourteen 28-day cycles) unless your doctor tells you to stop. Your doctor will monitor you while on treatment, and will stop treatment if your disease progresses or if you develop certain side effects.

Important Safety Information About CALQUENCE® (acalabrutinib) (cont'd)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking CALQUENCE with certain other medications may affect how CALQUENCE works and can cause side effects. Especially tell your healthcare provider if you take a blood thinner medicine.

How should I take CALQUENCE?

- Take CALQUENCE exactly as your healthcare provider tells you to take it.
- Do not change your dose or stop taking CALQUENCE unless your healthcare provider tells you to.
- Your healthcare provider may tell you to decrease your dose, temporarily stop, or completely stop taking CALQUENCE if you develop certain side effects.
- Take CALQUENCE 2 times a day (about 12 hours apart).
- Take CALQUENCE with or without food.

CALQUENCE + VENETOCLAX FIXED-DURATION TREATMENT

FOR PREVIOUSLY UNTREATED CLL/SLL

How CALQUENCE + venetoclax works



CALQUENCE blocks a protein called BTK to help stop the growth of CLL/SLL cells.*



Venetoclax works differently to destroy CLL/SLL cells by blocking a protein called B-cell lymphoma-2.*

*Healthy cells may also be affected.

How to take CALQUENCE + venetoclax[†]

	Cycles 1-2	Cycle 3	Cycles 4-14 [‡]
CALQUENCE	 1 x 100-mg tablet twice a day [§]	 1 x 100-mg tablet twice a day [§]	 1 x 100-mg tablet twice a day [§]
venetoclax		Week 1  2 x 10-mg tablets daily Week 2  1 x 50-mg tablet daily Week 3  1 x 100-mg tablet daily Week 4  2 x 100-mg tablets daily	 4 x 100-mg tablets daily Tablets not actual size.

Treatment cycles (each cycle is 28 days, or 4 weeks)

[†]Take each tablet by mouth.

[‡]Stop treatment after Cycle 14 unless your doctor tells you to stop earlier because your disease progresses or you develop certain side effects.

[§]About 12 hours apart as prescribed by your doctor.



It is important to hydrate before and during your treatment with venetoclax. Please talk to your doctor for more information.

Important Safety Information About CALQUENCE® (acalabrutinib) (cont'd)

- Swallow CALQUENCE tablets whole with a glass of water. Do not chew, crush, dissolve, or cut tablets.
- If you miss a dose of CALQUENCE, take it as soon as you remember. If it is more than 3 hours past


your usual dosing time, skip the missed dose and take your next dose of CALQUENCE at your regularly scheduled time. Do not take an extra dose to make up for a missed dose.

Please see additional Important Safety Information throughout and click here for full [Prescribing Information](#), including [Patient Information](#).


CALQUENCE®
acalabrutinib 100 mg tablets

CALQUENCE CONTINUOUS TREATMENT

FOR PREVIOUSLY UNTREATED AND PREVIOUSLY TREATED CLL/SLL



How CALQUENCE works



CALQUENCE blocks a protein called BTK to help stop the growth of CLL/SLL cells.*

*Healthy cells may also be affected.

How to take CALQUENCE

CALQUENCE



1 x 100-mg tablet
twice a day^{†‡}

Take CALQUENCE until your doctor tells you to stop.

[†]Take each tablet by mouth.

[‡]About 12 hours apart as prescribed by your doctor.



DISCOVER MORE

Important Safety Information About CALQUENCE® (acalabrutinib) (cont'd)

What are the possible side effects of CALQUENCE?

CALQUENCE can cause serious side effects, including:

- **Serious infections** have happened during treatment with CALQUENCE and may lead to death. Your healthcare provider may prescribe certain medicines if you have an increased risk of getting infections. Tell your healthcare provider right away if you have any signs or symptoms of an infection, including fever, chills, or flu-like symptoms.
- **Bleeding problems (hemorrhage)** have happened in people treated with CALQUENCE and can be serious and may lead to death. Your risk of bleeding may increase if you are also taking a blood thinner medicine. Tell your healthcare provider right away if you have any signs or symptoms of bleeding, including blood in your stools or black stools (looks

CALQUENCE INSTRUCTIONS & TIPS

FOR ALL AVAILABLE TREATMENT OPTIONS

Instructions for taking CALQUENCE



Take one 100-mg tablet by mouth, approximately every 12 hours



Take with or without food



Swallow tablet whole with a glass of water. Do not chew, crush, dissolve, or cut tablets

Tips for taking CALQUENCE

- Do not change your dose or stop taking CALQUENCE unless your doctor tells you to
- Your doctor may tell you to decrease your dose, temporarily stop, or completely stop taking CALQUENCE if you develop certain side effects
- If you miss a dose of CALQUENCE, take it as soon as you remember. If it's more than 3 hours past your usual dosing time, skip the missed dose and take your next dose of CALQUENCE at your regularly scheduled time. Do not take an extra dose to make up for a missed dose

Take CALQUENCE as directed by your doctor.



LEARN MORE

Important Safety Information About CALQUENCE® (acalabrutinib) (cont'd)

like tar), pink or brown urine, unexpected bleeding or bleeding that is severe or you cannot control, vomit blood or vomit that looks like coffee grounds, cough up blood or blood clots, dizziness, weakness, confusion, changes in your speech, headache that lasts a long time, or bruising or red or purple skin marks.

- **Decrease in blood cell counts.** Decreased white blood cell counts, red blood cell counts, and platelet counts can be severe during treatment with CALQUENCE. Your healthcare provider should do blood tests to

check your blood counts regularly during treatment with CALQUENCE.

- **Second primary cancers.** New cancers have happened in people during treatment with CALQUENCE, including cancers of the skin or other organs. Your healthcare provider will check you for skin cancers during treatment with CALQUENCE. Use sun protection when you are outside in sunlight.

Please see additional Important Safety Information throughout and click here for full [Prescribing Information](#), including [Patient Information](#).


CALQUENCE
acalabrutinib 100 mg tablets

HOW TO STAY ON TRACK



Not an actual patient.

The power of consistency

It's important to follow your doctor's treatment instructions and not miss doses. Here are some helpful tips to remember to take your treatment regimen.



Set reminders

Use alarms or a calendar to keep track of dosing times



Stay organized

Use a pillbox to ensure you take the right number of tablets each day



Keep a routine

Take your medicine during a routine like brushing your teeth



Don't forget

Check that you pack the right number of doses when traveling

Important Safety Information About CALQUENCE® (acalabrutinib) (cont'd)

What are the possible side effects of CALQUENCE? (cont'd)

CALQUENCE can cause serious side effects, including: (cont'd)

- **Heart rhythm problems (cardiac arrhythmias)** have happened in people treated with CALQUENCE, and can be serious or lead to death. Tell your healthcare provider right away if you have any of

the following signs or symptoms: fast or irregular heartbeat, dizziness, feeling faint, chest discomfort, or shortness of breath.

- **Liver problems.** Liver problems have happened in people treated with CALQUENCE, and can be severe or life-threatening, or lead to death. Tell your healthcare provider right away if you experience stomach pain or discomfort, urine of dark color or yellowing of your skin. Your healthcare provider will request tests to monitor your liver function during treatment with CALQUENCE.

POSSIBLE SIDE EFFECTS

Some side effects may be managed at home

Below are some of the most common side effects of CALQUENCE and tips on how to manage them.



Diarrhea

- Avoid caffeine, alcohol, dairy, fat, fiber, and spicy foods
- Eat low-fiber foods
- Talk to your doctor about taking an anti-diarrheal medication (like Imodium®)



Muscle and joint pain

- Take acetaminophen (Tylenol®)
- Discuss therapeutic massage or physical therapy with your doctor
- Apply heat or cold to the affected area



Upper respiratory tract infection

- Talk to your doctor right away if you have any signs or symptoms of an infection, including fever, chills, or flu-like symptoms
- Wash hands with soap and water
- Wear a mask when in crowds



Headache

- Take acetaminophen (Tylenol®)
- Get 7-9 hours of sleep
- Caffeine (coffee or tea)

These are not all the possible side effects of CALQUENCE.

Call your doctor if you experience any side effects and discuss how to manage them.

You may report side effects to FDA at 1-800-FDA-1088.



SEE MORE TIPS

Important Safety Information About CALQUENCE® (acalabrutinib) (cont'd)

The most common side effects of CALQUENCE

include upper respiratory infection, diarrhea, headache, muscle and joint pain, decreased white blood cell counts, and decreased platelet counts.

What is CALQUENCE?

CALQUENCE is a prescription medicine used to treat adults with:

- Chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL).

It is not known if CALQUENCE is safe and effective in children.

Please see additional Important Safety Information throughout and click here for full Prescribing Information, including Patient Information.


CALQUENCE®
acalabrutinib 100 mg tablets

RESOURCES



Not an actual patient and caregiver.

The power of support

The CALQUENCE™ Cares program offers resources and support during the different phases of your CLL/SLL treatment, including a free starter kit.



SIGN UP

Peer support

The CALQUENCE™ Connections Facebook Group is a safe and supportive online community to connect with and learn from others living with CLL/SLL.



JOIN TODAY



MOBILE TOOLS



The power of connection

Taking your treatment as prescribed and connecting with others who have similar cares and concerns as you can help better your treatment journey with CLL/SLL.

KEEP TRACK



Medisafe

The #1 medication reminder app, according to pharmacists surveyed



MyTherapy

Specifically developed for those with long-term health conditions

STAY CONNECTED



BELONG Beating Cancer Together

A social network for navigating cancer treatment by connecting patients, caregivers, and leading doctors

AstraZeneca provides these links as a service to you, and is not responsible for the content or privacy policy of any third-party app you choose. We encourage you to read the privacy policy of every application you choose.



EXPLORE RESOURCES

You may **report side effects related to AstraZeneca products** [↗](#).

Please see additional Important Safety Information throughout and **[click here for full Prescribing Information, including Patient Information.](#)**


CALQUENCE[®]
acalabrutinib 100 mg tablets

Important Safety Information About CALQUENCE® (acalabrutinib)

Before taking CALQUENCE, tell your healthcare provider about all of your medical conditions, including if you:

- have had recent surgery or plan to have surgery. Your healthcare provider may stop CALQUENCE for any planned medical, surgical, or dental procedure.
- have bleeding problems.
- have or had heart rhythm problems.
- have an infection.
- have or had liver problems, including hepatitis B virus (HBV) infection.
- are pregnant or plan to become pregnant. CALQUENCE may harm your unborn baby and cause problems during childbirth (dystocia).

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with CALQUENCE
- Use effective birth control (contraception) during treatment with CALQUENCE and for 1 week after the last dose of CALQUENCE
- are breastfeeding or plan to breastfeed. It is not known if CALQUENCE passes into your breast milk. Do not breastfeed during treatment with CALQUENCE and for 2 weeks after your last dose of CALQUENCE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking CALQUENCE with certain other medications may affect how CALQUENCE works and can cause side effects. Especially tell your healthcare provider if you take a blood thinner medicine.

How should I take CALQUENCE?

- Take CALQUENCE exactly as your healthcare provider tells you to take it.
- Do not change your dose or stop taking CALQUENCE unless your healthcare provider tells you to.
- Your healthcare provider may tell you to decrease your dose, temporarily stop, or completely stop taking CALQUENCE if you develop certain side effects.
- Take CALQUENCE 2 times a day (about 12 hours apart).
- Take CALQUENCE with or without food.
- Swallow CALQUENCE tablets whole with a glass of water. Do not chew, crush, dissolve, or cut tablets.
- If you miss a dose of CALQUENCE, take it as soon as you remember. If it is more than 3 hours past your usual dosing time, skip the missed dose and take your next dose of CALQUENCE at your regularly scheduled time. Do not take an extra dose to make up for a missed dose.

What are the possible side effects of CALQUENCE?

CALQUENCE can cause serious side effects, including:

- **Serious infections** have happened during treatment with CALQUENCE and may lead to death. Your healthcare provider may prescribe certain medicines if you have an increased risk of getting infections. Tell your healthcare provider right away if you have any signs or symptoms of an infection, including fever, chills, or flu-like symptoms.
- **Bleeding problems (hemorrhage)** have happened in people treated with CALQUENCE and can be serious and may lead to death. Your risk of bleeding may increase if you are also taking a blood thinner medicine. Tell your healthcare provider right away if you have any signs or symptoms of bleeding, including blood in your stools or black stools (looks like tar), pink or brown urine, unexpected bleeding or bleeding that is severe or you cannot control, vomit blood or vomit that looks like coffee grounds, cough up blood or blood clots, dizziness, weakness, confusion, changes in your speech, headache that lasts a long time, or bruising or red or purple skin marks.
- **Decrease in blood cell counts.** Decreased white blood cell counts, red blood cell counts, and platelet counts can be severe during treatment with CALQUENCE. Your healthcare provider should do blood tests to check your blood counts regularly during treatment with CALQUENCE.
- **Second primary cancers.** New cancers have happened in people during treatment with CALQUENCE, including cancers of the skin or other organs. Your healthcare provider will check you for skin cancers during treatment with CALQUENCE. Use sun protection when you are outside in sunlight.
- **Heart rhythm problems (cardiac arrhythmias)** have happened in people treated with CALQUENCE, and can be serious or lead to death. Tell your healthcare provider right away if you have any of the following signs or symptoms: fast or irregular heartbeat, dizziness, feeling faint, chest discomfort, or shortness of breath.
- **Liver problems.** Liver problems have happened in people treated with CALQUENCE, and can be severe or life-threatening, or lead to death. Tell your healthcare provider right away if you experience stomach pain or discomfort, urine of dark color or yellowing of your skin. Your healthcare provider will request tests to monitor your liver function during treatment with CALQUENCE.

The most common side effects of CALQUENCE include upper respiratory infection, diarrhea, headache, muscle and joint pain, decreased white blood cell counts, and decreased platelet counts.

These are not all the possible side effects of CALQUENCE. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

What is CALQUENCE?

CALQUENCE is a prescription medicine used to treat adults with:

- Chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL).

It is not known if CALQUENCE is safe and effective in children.

You may report side effects related to AstraZeneca products .

Please see additional Important Safety Information throughout and click here for full Prescribing Information, including Patient Information.


CALQUENCE[®]
acalabrutinib 100 mg tablets

FINANCIAL SUPPORT

More than 90% of patients prescribed CALQUENCE pay \$0 out of pocket, regardless of the type of insurance coverage.

FOR COMMERCIALLY INSURED PATIENTS



CALQUENCE Co-pay Savings Program

Patients may pay as little as \$0 per month for CALQUENCE. There are no income requirements to participate in the program. For more information and eligibility requirements, visit www.azpatientsupport.com.

*Terms and conditions apply. See site for full eligibility and terms of use.

FOR GOVERNMENT-INSURED OR UNINSURED PATIENTS



Independent Foundations

AstraZeneca Access 360™ can provide information about independent charitable patient assistance foundations that may be able to help you with out-of-pocket costs. However, Access 360 does not guarantee support by independent foundations.

For more information and a list of foundations, call Access 360 at 1-844-ASKA360 or visit <https://www.myaccess360.com/patient/calquence-acalabrutinib.html>.



AZ&ME Prescription Savings Program

AZ&ME provides AstraZeneca medicines at no cost to qualifying people. For more information, call 1-800-AZandMe (1-800-292-6363) or visit www.AZandMeApp.com.

MEDICARE PART D



Learn more about the 2026 Medicare Part D Reform updates. Visit <https://www.azmedicareupdates.com/>.



CALQUENCE is a registered trademark, and AstraZeneca Access 360 and AZ&Me are trademarks of the AstraZeneca group of companies.
©2026 AstraZeneca. All rights reserved. US-108008 Last Updated 4/26

